

The co-evolution of the obscured quasar PKS1549-79 and its host galaxy: evidence for a high accretion rate and warm outflow

J. Holt¹, C. Tadhunter¹, R. Morganti^{2,3}, M. Bellamy¹, R.M. González Delgado⁴, A. Tzioumis⁵, K.J. Inskip¹

¹*Department of Physics and Astronomy, University of Sheffield, Sheffield, S3 7RH, UK*

²*ASTRON, PO Box 2, 7990 AA Dwingeloo, The Netherlands*

³*Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands*

⁴*Instituto de Astrofísica de Andalucía, Apdo. 3004, 18080 Granada, Spain*

⁵*ATNF-CSIRO, Epping, Sydney, Australia*

ABSTRACT

We use deep optical, infrared and radio observations to explore the symbiosis between the nuclear activity and galaxy evolution in the southern compact radio source PKS1549-79 ($z = 0.1523$). The optical imaging observations reveal the presence of tidal tail features which provide strong evidence that the host galaxy has undergone a major merger in the recent past. The merger hypothesis is further supported by the detection of a young stellar population, which, on the basis of spectral synthesis modelling of our deep VLT optical spectra, was formed 50 – 250 Myr ago and makes up a significant fraction of the total stellar mass (1 – 30%). Despite the core-jet structure of the radio source, which is consistent with the idea that the jet is pointing close to our line of sight, our HI 21cm observations reveal significant HI absorption associated with both the core and the jet. Moreover the luminous, quasar-like AGN ($M_V < -23.5$) is highly extinguished ($A_v > 4.9$) at optical wavelengths and shows many properties in common with narrow line Seyfert 1 galaxies (NLS1), including relatively narrow permitted lines ($FWHM \sim 1940 \text{ km s}^{-1}$), highly blueshifted [OIII] $\lambda\lambda 5007, 4959$ lines ($\Delta V \sim 680 \text{ km s}^{-1}$), and evidence that the putative supermassive black hole is accreting at a high Eddington ratio ($0.3 < L_{bol}/L_{edd} < 11$). The results suggest that accretion at high Eddington ratio does not prevent the formation of powerful relativistic jets.

Together, the observations lend strong support to the predictions of some recent numerical simulations of galaxy mergers in which the black hole grows rapidly through merger-induced accretion following the coalescence of the nuclei of two merging galaxies, and the major growth phase is largely hidden at optical wavelengths by the natal gas and dust. Although the models also predict that AGN-driven outflows will eventually remove the gas from the bulge of the host galaxy, the *visible* warm outflow in PKS1549-79 is not currently capable of doing so. However, much of the outflow may be hidden by the material obscuring the quasar and/or tied up in hotter or cooler phases of the interstellar medium.

By combining our estimates of the reddening of the quasar with the HI column derived from the 21cm radio observations we have also made the first direct estimate of the HI spin temperature in the vicinity of a luminous AGN: $T_{spin} > 3000\text{K}$.

Key words: galaxies:starburst – galaxies:individual: PKS1549-79 – galaxies:formation – galaxies:interactions – quasars:general

1 INTRODUCTION

Powerful radio galaxies have long been considered as useful signposts to massive structures in the distant universe.

However, there is now increasing speculation that AGN and associated jet activity are not merely signposts but, in fact, intimately linked to the galaxy evolution process. The main evidence for such links is twofold. On the one hand the shape of the redshift evolution in the number density of quasars and radio galaxies shows marked similarities to that of the star formation history of the universe (Dunlop & Peacock 1990, Madau et al. 1996), while on the other close correlations have been found between the masses of the supermassive black holes that produce the prodigious nuclear activity and the properties of the bulges of the host galaxies (e.g. Tremaine et al. 2002).

The correlations between the redshift evolution of the radio source population and the star formation history of the field galaxy population can be explained in terms of hierarchical galaxy evolution models (e.g. Kauffmann & Haenelt 2000). It is well known from studies of nearby merging galaxies that stars are formed as part of the merger process. Numerical simulations of mergers of gas rich galaxies also show that the tidal torques associated with the mergers concentrate gas in the circumnuclear regions of the galaxies (Barnes & Hearnquist 1996, Mihos & Hearnquist 1996); some of this gas may be available to fuel circumnuclear starburst, quasar and jet activity. Therefore it is likely that AGN activity and near-nuclear starbursts in massive galaxies are a direct consequence of hierarchical galaxy evolution; AGN may be a symptom of the overall galaxy evolution process.

In this context, radio galaxies are key objects for understanding the link between star formation and AGN activity in individual galaxies. Not only is there clear morphological evidence that a significant subset of powerful radio galaxies — $\sim 50\%$ of those with strong emission lines — have undergone recent mergers (Heckman et al. 1996, Smith & Heckman 1989), but many of the radio galaxies with morphological evidence for mergers also show signs of recent star formation (Tadhunter et al. 1996, Aretxaga et al. 2001, Tadhunter et al. 2002, 2005, Wills et al. 2002, 2004, Johnston et al. 2005). Despite these results, considerable uncertainties remain about the relationship between star formation and AGN/jet activity in radio galaxies. For example, it is not certain at what stage of the merger the AGN activity is triggered, nor whether the star formation is exactly coeval with the AGN activity. Indeed, some recent studies of the young stellar populations in radio galaxies suggest that the activity may be triggered relatively late in the merger sequence, a significant time after the major episode of merger-induced star formation (Tadhunter et al. 2005, Emonts et al. 2005, Johnston et al. 2005). However, the sample sizes are currently small and further detailed studies of the young stellar populations in powerful radio galaxies are required to put our understanding of the link between star formation and AGN activity on a firmer footing.

While the links between star formation and AGN/jet activity provide evidence that AGN are a bi-product of the overall hierarchical galaxy evolution process, it has been suggested that the correlations between the supermassive black holes masses and the properties of the bulges of the host galaxies are a consequence of the feedback effects of AGN-induced outflows (Silk & Rees 1998, Fabian 1999, di Matteo et al. 2005). To date, most of the evidence for such outflows has been provided by observations of narrow and broad absorption lines observed in the spectra of AGN at UV and

X-ray wavelengths (see Crenshaw, Kraemer & George 2003 for a review). The outflows associated with these absorption features clearly have an impact on the near-nuclear environments on a scale $< 10\text{pc}$, but the impact they have on the $1 - 10\text{ kpc}$ scale of galaxy bulges is less clear. Although outflows have also been detected in both ionized and neutral gas on a kpc-scale in powerful radio galaxies, uncertainties remain about the general importance of the outflows and their driving mechanism (quasars, jets or starbursts).

The subject of this paper — PKS1549-79 — is a key object for studies of the links between galaxy evolution and AGN activity, because it shows clear evidence of both the merging process that may trigger the activity, and the feedback effect of the AGN activity on the surrounding ISM.

PKS1549-79 is a compact flat spectrum radio source. Previously unpublished VLBI radio images of the source show that it has the compact core-jet structure expected for such sources, consistent with the idea that its jet is pointing close to the line of sight. Remarkably for such a source, low-resolution optical spectra fail to reveal the broad permitted lines and strong non-stellar continuum characteristic of quasars and BL Lac objects (Tadhunter et al. 2001). Therefore this object does not readily fit in with the simplest versions of the unified schemes which propose that lines of sight close to the radio axis have relatively unobscured views of the AGN (Barthel 1989).

Despite the absence of strong quasar continuum and broad emission line features, optical spectra of the source do reveal high ionization forbidden emission lines which suggest that the source contains a powerful quasar-like AGN, even if this AGN is not directly visible at optical wavelengths. However, the high ionization emission lines are unusually broad, and blueshifted by $\sim 650\text{ km/s}$ relative to the low ionization lines. In addition, this object has HI 21cm absorption detected against its radio core, at the same redshift as the low ionization emission line system. Further support for the idea that the quasar nucleus is obscured by the circumnuclear material in this source is provided by K-band spectroscopic observations that detect a broad $\text{Pa}\alpha$ emission line and strong non-stellar continuum characteristic of quasars (Bellamy et al. 2003).

Aside from its unusual emission line properties, PKS1549-79 belongs to the subset of $\sim 30 - 50\%$ of powerful radio galaxies for which young stellar populations make a substantial contribution to the optical continuum emission (Tadhunter et al. 2002). Further evidence for star formation in this object is provided by its unusually strong far-IR emission, which leads to its classification as an ultra luminous infrared galaxy (ULIRG: see Sanders & Mirabel 1996 for a definition) with $L_{\text{ir}} = 1.6 \times 10^{12} L_{\odot}$. This evidence for recent star formation is consistent with the idea that the host galaxy of this source has been involved in a merger which has also triggered a major episode of star formation.

In this paper we present new deep spectroscopic and imaging observations of PKS1549-79 at optical, infrared and radio wavelengths that allow us to investigate the triggering of its AGN activity, characterise its circumnuclear outflows, and deduce the nature of its AGN. In this way we can place this key object in the context of evolutionary scenarios for the origin and evolution of the activity.

We assume cosmological parameters of $H_0 = 75\text{ km s}^{-1}$ and $q_0 = 0.0$ throughout the paper. For these parameters

1 arcsec corresponds to 2.39 kpc at the redshift of the host galaxy of PKS1549-79.

2 OBSERVATIONS AND REDUCTIONS

2.1 Optical spectroscopy

Optical spectroscopic observations were taken in 2002 July and 2003 September using the EMMI spectrograph in RILD mode on the ESO 3.5-m New Technology Telescope (NTT), and the FORS2 spectrograph in LSS mode on the ESO Very Large Telescope (VLT). Both sets of data are deeper and have a higher spectral resolution than those reported in Tadhunter et al. (2001). Details of the observations are given in Table 1.

On the NTT, grisms #5 and #6 were used with the MIT/LL mosaic to obtain spectra with wavelength ranges 3215–6136 Å and 4977–7565 Å in the blue and red respectively. These configurations resulted in spectral resolutions of 5.0 – 6.7 Å in the blue and 4.2 – 5.6 Å in the red, depending on slit width. Red and blue exposures were interleaved in an attempt to ensure uniformity in conditions between the exposures. To reduce the effects of differential refraction, all exposures were taken with the slit aligned along the parallactic angle. Spectra were taken along two slit positions – PA 25 and PA -5.

The aim of the NTT spectra was to investigate the kinematics and physical conditions of the emission line gas through modelling of the bright emission lines. However, higher signal-to-noise is required to investigate the stellar populations in PKS 1549-79. Hence, we have also obtained optical spectra with the FORS2 spectrograph on the VLT on 2003 September 25. Grisms 600B and 600RI (holographic) were used with the MIT CCD to obtain spectra with wavelength ranges 3100–6000 Å and 5000–8200 Å respectively. The VLT spectra were taken with a 1.3 arcsec slit to ensure the entire nuclear region was sampled, and the slit was aligned along the parallactic angle (PA 75). The resulting spectral resolutions were 6.5 Å and 7.4 Å for the blue and red spectra respectively. As with the NTT observations, the red and blue exposures were interleaved.

For both the NTT and the VLT observations the data reduction followed the standard steps of bias subtraction, flat fielding, wavelength calibration, atmospheric extinction correction, flux calibration, and registration of blue and red spectral images. Based on measurements of night sky lines, the uncertainty in the wavelength calibration is estimated to be smaller than 0.5 Å for all datasets; the relative flux calibration uncertainty, based on the comparison of observations of several spectroscopic standard stars, is better than $\pm 5\%$. Prior to the analysis of the data, the spectra were corrected for a Galactic extinction assuming $E(B - V) = 0.679$ (Schlegel, Finkbeiner & Davis 1998) and the Seaton (1979) extinction law.

2.2 Optical imaging

Deep images of PKS1549-79 were obtained using FORS1 on the VLT with a Gunn r filter in good seeing conditions ($FWHM = 0.67$ arcseconds). The advantage of this filter is that, covering the rest wavelength range $\sim 5200 - 6100$ Å, it

Date	Exposure (s)	PA	Slit width (arcsec)	Seeing FWHM (arcsec)
NTT/EMMI (Optical Spectroscopy)				
12/07/02	3*1200(R)	-5	1.0	0.78
	3*1200(B)	-5	1.0	0.78
13/07/02	2*1200(R)	25	1.5	0.78
	2*1200(B)	25	1.5	0.78
VLT/FORS1 (Gunn r Imaging)				
25/07/03	4*300	-	-	0.67
VLT/FORS2 (Optical Spectroscopy)				
25/09/03	3*600(R)	75	1.3	1.2-1.8
25/09/03	3*1200(B)	75	1.3	1.2-1.8
AAT/IRIS2 (K-band Spectroscopy)				
4/09/03	24*120	0	1.0	1.0
NTT/SOFI (H and K-band Spectroscopy)				
24/08/04	12*120	0	1.0	1.2

Table 1. Log of optical and infrared observations for PKS1549-79. In the case of the optical spectroscopic observations the seeing was estimated from the DIMM monitor estimates at the time of the observations, whereas for the imaging observations the seeing was measured directly from star images in the field of the source.

avoids all strong emission lines in the source and is therefore dominated by continuum light. During the observations the telescope was jittered to four separate positions on the sky to help remove cosmetic defects and average out sensitivity variations.

The imaging data were reduced using a combination of the STARLINK FIGARO package and IRAF, by first subtracting the bias, dividing each separate frame by a jittered sky flat field, then combining the images with IMCOMBINE in IRAF using the coordinate information in the image headers.

2.3 Near-IR spectroscopy

In an attempt to improve on the near-IR spectroscopic data of Bellamy et al. (2003), which were affected by a faulty grism in IRIS2, new near-IR spectra of PKS 1549-79 were taken with the Anglo Australian Telescope (AAT), and the NTT. Details of the observations are given in Table 1.

The first data set was obtained on 2003 September 4 using the IRIS2 instrument on the AAT. The Sapphire 240 grism was used to cover the K band. In total, twenty four exposures of 120s were obtained for PKS 1549-79, with the galaxy ‘nodded’ between two positions on the slit in an ABBA pattern. Four exposures of 66s were taken of the A0 star HIP 77941 at a similar airmass for calibration purposes.

The second set was obtained on 2004 August 24 with the SOFI instrument on the NTT. The low resolution red grism was used with the GRF order-sorting filter to cover the H and K bands simultaneously. Cloud was present during the night and the observations were not photometric. 12 exposures of 120s were obtained; the galaxy was again nodded along the slit but this time with a random jitter every 2 exposures. Eight exposures of 30s were taken of the A0 star HIP 79933 at a similar airmass for calibration purposes.

The IRIS2 exposures were co-added using median filtering to remove cosmic rays; the median ‘B’ image was then subtracted from the median ‘A’ image to remove the night sky lines. The standard star frames were combined in a similar way. The resulting galaxy and star frames were then flat-fielded using a dark-subtracted flat-field taken with the same instrumental setup. The night sky lines from the data frames themselves were used to determine a two-dimensional wavelength calibration, which was then applied to the final co-added frames of the galaxy and the standard star. The resulting spectra cover the range 20320 – 24900 Å. The spectral resolution, measured from the night sky lines, is 10.9 ± 0.4 Å(FWHM). Fits to the night sky lines show that the wavelength calibration is accurate to ± 0.16 Å. The two-dimensional spectra were straightened using the APSUM package in IRAF, and one-dimensional spectra were extracted with FIGARO.

Night sky lines were also used to derive a wavelength calibration for the NTT data. The resulting two-dimensional transformation was then applied to all the individual data frames and then suitable groups of exposures (because of the random jittering) were combined with FIGARO and a series of one-dimensional spectra were extracted and then combined together. There was no appreciable tilting of the slit image and so the SOFI spectra did not need to be straightened before extraction. The SOFI spectrum covers the wavelength range 15057 – 25392 Å. Atmospheric absorption dominates between ~ 18000 and 19500 Å and the thermal background dominates at wavelengths greater than ~ 24000 Å. The spectral resolution, measured from the arc lines, is 35.0 ± 0.4 Å(FWHM). Fits to the night sky lines show that the wavelength calibration is accurate to ± 0.77 Å.

The standard star exposures were used to remove telluric features and also to flux calibrate the IRIS2 spectra. The magnitude-to-flux conversion was performed with reference to Bessell, Castelli & Plez (1998), while telluric correction (and flux calibration for the IRIS2 data) was performed by assuming the intrinsic SED of the A0 stars to be that of a black body at $T = 9480$ K. For hot stars this is a good approximation, particularly in the near-IR end of the black body spectrum. A number of Brackett lines are visible in the spectra of A0 stars, particularly in the H band, but these were masked out before the telluric correction was performed.

The SOFI data suffered unquantifiable slit-losses due to the telescope nodding axis not being aligned precisely with the slit. For this reason no attempt was made to make an absolute flux calibration of the SOFI spectrum. Instead the spectrum was scaled to match the IRIS2 spectrum in the K-band and the spectrum of HIP 79933 was used only to remove telluric features and obtain the correct spectral shape.

2.4 Radio observations

Continuum VLBI observations of PKS1549-79 were made at both 2.3 and 8.4 GHz with the SHEVE network (Preston et al. 1993, Jauncey et al. 1994) between November 1988 and March 1992, using the MK2 recording system (Clark 1973). Full details of the antennae used for SHEVE observations are given in Tzioumis et al. (2002). The tapes recorded at

each station were processed at the Caltech-JPL Block2 correlator in Pasadena, and the NRAO AIPS package was used for global fringe fitting. The amplitude calibration and editing of the visibilities were subsequently performed using the Caltech VLBI package. Self-calibration and imaging of the data were performed using DIFMAP (Shepherd et al. 1994) and the AIPS packages. At 2.3 GHz the restoring beams were 7.3×2.7 mas (PA4.7) and 8.9×7.2 mas (PA9.8) with and without the Harteebeesthoek antenna respectively, whereas at 8.4 GHz the restoring beam was 6.7×5.2 mas (PA73) without the Harteebeesthoek antenna. The r.m.s. noise in the final images is 2.7 mJy beam $^{-1}$ at 2.3 GHz, and 3.8 mJy beam $^{-1}$ at 8.4 GHz.

In order to investigate the spatial distribution of the HI 21cm absorption previously detected by Morganti et al. (2001), HI VLBI observations were carried out using the Long Baseline Array (LBA) at the redshifted HI frequency of 1645 MHz. At this relatively low frequency we could use only three telescopes: Parkes (64m), ATCA (Tied Array, 5×22 m) and Mopra (22m). We used a 16 MHz bandwidth in each circular polarization and 1024 channels. The final data cube (after Hanning smoothing) has a spectral resolution of ~ 9 km s $^{-1}$, and a noise per spectral channel of 5.3 mJy beam $^{-1}$, while the continuum map derived from these data has a noise of 16 mJy beam $^{-1}$.

The editing part of the calibration of the 21cm line data was done in AIPS and then the data were transferred to MIRIAD (Sault, Teuben & Wright 1995) for the bandpass calibration. The calibration of the bandpass was done using additional calibrations of the strong calibrators PKS1718-649 and PKS1934-638.

Both the line and the continuum image were made using uniform weighting and the restoring beam is 47×32 mas (PA-77.3). As expected, the resolution is lower than that obtained in the previous VLBI continuum observations described above. The total cleaned flux in the continuum image is 4.75 Jy — close to that obtained from the previous low resolution ATCA observations.

3 RESULTS

3.1 Optical imaging

Our deep VLT image of PKS1549-79 is shown in Figure 1. This reveals a disturbed optical continuum morphology, with a high surface brightness elongation extending 4 arcseconds (10 kpc) to the north of the nucleus, a pair of tidal tails (or loop feature observed close to edge-on) extending a total distance of 10 arcseconds (24 kpc) to the south, and a fainter jet-like feature that extends 12 arcseconds (29 kpc) to the NNE of the nucleus. The extended structures contain several knots of emission, particularly to the SSE and NNW. Overall, the structures suggest that PKS1549-79 has undergone a recent merger, which induced star formation in the extended halo of the galaxy.

3.2 Optical spectroscopy

Three slit positions used for the optical spectroscopy observations are shown superimposed on the Gunn r image in Figure 1.




Figure 1. Deep VLT image of PKS1549-79 taken with a Gunn r filter that avoids strong emission lines from the galaxy. The slit positions used for the long-slit spectroscopic observations are shown for reference. North is to the top and east to the left.

3.2.1 Optical continuum modelling

Accurate modelling of the continuum is required not only to deduce the properties of any young stellar populations present, but also as a prerequisite for accurate measurement of the emission lines, following subtraction of the best-fit continuum model. In this section we concentrate on the VLT spectrum extracted for the nuclear region because this has higher S/N in the continuum than either of the NTT spectra.

Prior to continuum modelling, the level of the nebular continuum was deduced following the procedure outlined in Dickson et al. (1995) and Holt et al. (2003), based on double Gaussian fits to the $H\beta$ emission line feature. This nebular continuum was then subtracted from the data. The major uncertainty in this procedure is the reddening in the nebular continuum. Unfortunately, estimates of the reddening using the Balmer lines are highly uncertain in this object (see section 3.2.3). To consider the effects of the reddening of the nebular continuum on the results of the continuum mod-

elling, we carried out the continuum modelling twice with two extreme sets of assumptions about nebular continuum. First we assumed zero reddening for the nebular continuum. Second we assumed the maximum reddening allowed by the Balmer line ratios.

The technique used to model the continuum is described in detail in Tadhunter et al. (2002, 2005). To summarise, the continuum flux is measured in several wavelength bins (in this case 39) chosen to avoid emission lines, cosmetic defects and poorly-corrected atmospheric absorption features. A minimum χ^2 technique is then used to compare the measured fluxes with the results of spectral synthesis models. Models comprising various combinations of an old stellar population (12.5 Gyr), young stellar populations (0.01 – 2.5 Gyr), reddened young stellar populations (0.01 – 2.5 Gyr, $0.0 < E(B - V) < 2.0$), and a power-law are compared with the data. For the spectral synthesis results we used the most recent set of high resolution models from Bruzual and Charlot (2003) for an instantaneous burst, a Salpeter IMF and

solar metallicity. For the reddened young stellar populations (YSP) we used the Seaton (1979) extinction law to redden the YSP spectra from the spectral synthesis models. We also assumed a uniform uncertainty of $\pm 5\%$ for all the measured fluxes, consistent with the estimated uncertainty of the relative flux calibration. Based on our experience of this modelling technique, models with $\chi^2_{red} < 1$ can be deemed to provide a good fit to the data (see Tadhunter et al. 2005 for discussion).

In our first attempt to model the nebular-subtracted optical continuum, we considered the full wavelength range covered by the spectra. It became clear that no combination of old and young stellar populations (reddened or otherwise) could fit the relatively steep rise in the spectrum to the red at wavelengths longer than $\sim 6500\text{\AA}$, and that a red power-law is required to provide an adequate fit in that spectral range. Note that the steep rise to the red is not due to any problem with the reduction and calibration of the data, since it is also present in our NTT spectra. Figure 2 shows the best fitting model that comprises a 12.5 Gyr old stellar population, a 0.05 Gyr young stellar population reddened by $E(B - V) = 0.4$, and a power-law. The level of the power-law component is consistent with the extrapolation of the reddened quasar continuum detected at near-IR wavelengths (see section 3.3). Indeed the existence of the red power-law provides independent evidence for the partially extinguished quasar first detected by Bellamy et al. (2003).

In order to avoid the added uncertainty that the inclusion of a power-law introduces into the modelling of the continuum, and to deduce an accurate age for the YSP, we then excluded wavelengths $\lambda > 6000\text{\AA}$, from the model fits. In the case of maximum reddening for the subtracted nebular continuum, it is clear that the data are well fitted by models that include a combination of a 12.5 Gyr old stellar population and a YSP with age in the range 0.04 – 0.1 Gyr and reddening in the range $0.0 < E(B - V) < 0.75$, with a best fit obtained for a YSP age of 0.05 Gyr and reddening of $E(B - V) = 0.6$. The results appear to be robust in the sense that, if we assume that the nebular continuum suffers no reddening, the results are similar, except that the range of acceptable models extends to older ages for the YSP (0.05 – 0.25 Gyr) and the best fitting model has an age of 0.1 Gyr and a reddening of $E(B - V) \sim 0.5$. The results of the SED modelling for the restricted wavelength range are shown in Figure 3 for the zero reddening in the nebular continuum.

The next step in the analysis was to check the SED fitting results by determining how well the models fit the profiles of absorption lines that do not suffer from significant emission line contamination, namely the CaII K, the G-band, and MgI features. Reassuringly, the models that fit the general SED well also provide the best fits to the absorption lines. Figure 4 shows an example of a fit to the optical spectrum for a model with a YSP of age 0.1 Gyr and reddening $E(B - V) = 0.5$, assuming zero reddening for the nebular continuum.

To summarise: fits to the optical continuum require a young stellar population (YSP) with age in the range 0.04 – 0.25 Gyr and reddening $0.0 < E(B - V) < 0.8$ (depending on age) in order to provide an adequate fit to the optical SED and stellar absorption features. The YSP has a mass $2 \times 10^8 < M_{YSP} < 4 \times 10^9 M_\odot$ and makes up 1 – 30% of the total stellar mass in the slit (depending on the exact age

and reddening). Note the age deduced for the YSP is significantly smaller than the 1 Gyr determined by Tadhunter et al. (2002) using lower resolution data with a narrower wavelength range. Much of this difference is likely to be due to the fact that Tadhunter et al. (2002) did not correct their spectrum for Galactic extinction prior to the modelling.

3.2.2 Emission line kinematics

The long-slit spectra of PKS1549-79 provide useful information on the emission line kinematics in both the nuclear and off-nuclear emission line regions. The spectra were measured by using the DIPSO package to fit Gaussian profiles to the lines. The velocity widths derived from the fits were quadratically corrected for the instrumental profiles, and all line widths and radial velocity shifts were corrected to the rest frame of PKS1549-79.

The NTT spectra, which were obtained under good seeing conditions, provide the best information about the spatial extent and kinematics of the extended emission line gas. Figure 5 presents flux and radial velocity information derived from the fits to the emission line profiles. The H α emission line is detected across the full extent of the higher surface brightness continuum structures detected in the Gunn r continuum image shown in Figure 1, extending 10 arcseconds to the south and 5 arcseconds to the north of the nucleus. The emission line kinematics are relatively quiescent in the extended regions with small velocity amplitudes ($\Delta V < 150 \text{ km s}^{-1}$) and linewidths ($\text{FWHM} < 400 \text{ km s}^{-1}$) — consistent with a gravitational origin for the gas motions (Tadhunter et al. 1989, Baum et al. 1990).

In contrast to the extended regions, the kinematics in the nuclear region provide evidence for more extreme gas motions, as already noted by Tadhunter et al. (2001). In order to analyse the gas motions in the spatially unresolved nuclear regions we have adopted the approach used by Holt et al. (2003) for the radio galaxy PKS1345+12. This consists of fitting the minimum number of Gaussian components required model each line adequately. For this analysis we have concentrated on the VLT data which have a higher S/N in the nuclear regions. In the case of emission line doublets ([OIII] $\lambda\lambda 5007, 4959$, [OI] $\lambda\lambda 6300, 6363$, [NII] $\lambda\lambda 6548, 6584$, [SII] $\lambda\lambda 6717, 6731$) the separations, and, with the exception of the [SII] blend, the ratios of the lines were set by atomic physics, and it was assumed that both components of the doublets have the same kinematic components. We found that for all the low ionization lines [OII] $\lambda 3727$, H β , [OI] $\lambda\lambda 6300, 6363$, [NII] $\lambda\lambda 6548, 6584$, and [SII] $\lambda\lambda 6717, 6731$, a double Gaussian fit comprising narrow ($\text{FWHM} < 400 \text{ km s}^{-1}$) and intermediate width blueshifted components ($\text{FWHM} > 1000 \text{ km s}^{-1}$) provided an adequate fit to the emission line profiles. For these low ionization lines the narrow component is strongest, and the widths and redshifts of the narrow components are consistent within the uncertainties for all the narrow lines; they are also consistent with the redshift of the H α emission line detected in the extended regions (see Figure 5). The intermediate emission line component is relatively weak in the low ionization lines, and its width and redshift less well determined based on free double Gaussian fits.

Initial attempts to fit the [OIII] $\lambda\lambda 5007, 4959$ lines showed that it was possible to fit each of line of the dou-




Figure 2. Spectral synthesis model fit to the full wavelength range of the optical spectrum of PKS1549-79. The model comprises a 0.05 Gyr YSP reddened by $E(B - V) = 0.4$, a 12.5 Gyr old stellar population, and a power-law. Note that the power-law component is required to fit the steep rise in the continuum at the far red end of the spectrum. The differences between the model and the data are presented as a histogram at the bottom of the plot.




Figure 3. The results of minimum χ^2 fitting to the optical continuum of PKS1549-79 assuming the zero reddening for the nebular continuum, restricting the wavelength range of the fits to $\lambda_{rest} < 6000\text{\AA}$. The plot on the left gives the reduced χ^2 of the model fits for different combinations of YSP age and reddening, while the plot on the right gives the percentage contribution of the YSP component in the normalising bin used for the modelling (4720 – 4820 Å). Reduced χ^2 values less than unity are deemed acceptable.

blet adequately using a single Gaussian component that is both broad and blueshifted relative to the narrow component detected in the low ionization lines (see also Tadhunter et al. 2001). However, given that results obtained for [OIII] in apertures slightly offset from the nucleus show the clear presence of a narrow ($\text{FWHM} < 400 \text{ km s}^{-1}$) emission line component that is also detected in the low ionization lines, we have also fitted each component of [OIII] doublet in the nucleus using a double Gaussian model, with one of the Gaussian components constrained to have the velocity width and redshift of the narrow component detected in the low ionization lines. The resulting fit — comprising the narrow component and an intermediate, blueshifted component — is shown in Figure 6. The imperfections of this fit in the line

cores are to likely reflect the fact that a single Gaussian is only an approximation to the true profile of the intermediate, blueshifted component in the nucleus. Note that such imperfections would not be apparent in fits to the low ionization lines, because of the weakness of the broad components, and the dominance of the narrow components in such lines.

Following on from the double Gaussian fits to [OIII] we then attempted to fit all the emission lines in the nuclear regions with a double Gaussian model comprising two components:

- A narrow component ($\text{FWHM} = 383 \pm 15 \text{ km s}^{-1}$) with redshift $z = 0.15225$, measured from double Gaussian fits to the bright low ionisation lines.




Figure 4. Example fit to the optical continuum for the model including a YSP with age 0.1 Gyr and reddening $E(B - V) = 0.5$ (zero reddening for the nebular continuum). The CaII K, G-band and MgI absorption features are well fitted by this model, but the higher order Balmer lines are likely to suffer from residual emission line contamination.

- An intermediate component ($\text{FWHM} = 1282 \pm 25 \text{ km s}^{-1}$) blue shifted by $679 \pm 20 \text{ km s}^{-1}$ with respect to the narrow component, measured from double Gaussian fits to the [OIII] lines.

This model provides an adequate fit to all the lines in the nuclear emission spectrum, with the following notable exceptions:

- The $\text{H}\alpha + [\text{NII}]$ blend requires an additional broad $\text{H}\alpha$ emission line component to fit the wings of the blend. We find that we can obtain an adequate fit to the blend if we set the width and redshift of the broadest $\text{H}\alpha$ component to be the same as those deduced from single Gaussian fits to the broad $\text{Pa}\alpha$ line detected by Bellamy et al. (2003: $z = 0.15238$, $\text{FWHM} = 1940 \text{ km s}^{-1}$). The flux measured for the broadest $\text{H}\alpha$ component is $4.2 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$. The detection of this component provides further evidence that the reddened quasar nucleus is detected at the longer wavelength end of the optical window.

- The faint high ionization lines [NeV] $\lambda 3425$ and [FeVII] $\lambda 6087$ could be fitted adequately by the broad component alone, whereas the faint low ionization [NI] $\lambda 5200$ could be fitted by the narrow component alone. Presumably, in each of these cases the other component that would otherwise be detected in a double Gaussian fit is not detected because of the low S/N and uncertainties in the continuum subtraction.

The velocity shifts and widths of the various kinematic components detected in PKS1549-79 are compared in Figure 5 and Table 2. Not surprisingly, the intermediate and narrow components in the double Gaussian fits have similar redshifts and line widths to the high and low ionization lines

Component	Redshift	Line width(FWHM) (km s^{-1})
Nuclear narrow	0.15225 ± 0.00003	383 ± 15
Nuclear intermediate	0.14959 ± 0.00003	1282 ± 25
$\text{Pa}\alpha$ core	0.15238 ± 0.00006	1250 ± 70
$\text{Pa}\alpha$ broadest	0.1529 ± 0.0002	3525 ± 200
HI 21cm narrow	0.15195 ± 0.00006	80 ± 30

Table 2. Comparison of the redshifts and line widths of the various kinematic components detected in the nuclear regions of PKS1549-79. Results for nuclear narrow and intermediate components are based on double Gaussian fits to the optical emission lines detected in the VLT spectrum, while the $\text{Pa}\alpha$ results are based on double Gaussian fits to the $\text{Pa}\alpha$ feature detected in the new AAT K-band spectrum.

measured in single Gaussian fits to the lower resolution data of Tadhunter et al. (2001). It is notable that both the width and velocity shift of the nuclear narrow component are consistent with those deduced for the extended $\text{H}\alpha$ emission on both sides of the nucleus; the redshift of this component is also close to that measured for both the core of the broad $\text{Pa}\alpha$ line and the narrow HI absorption (see below). Therefore we identify the nuclear narrow component with the rest frame of the host galaxy, and the intermediate component detected most clearly in the [OIII] lines with a highly disturbed component that is blueshifted relative to this rest frame.




Figure 5. Radial velocity profiles (top) and the spatial variations of emission line flux (bottom) for the NTT observations of PKS 1549-79. The left hand images are for PA-5 and the right hand images are for PA25. Small open and filled triangles represent the narrow and intermediate components of H α respectively. Overplotted is the radial velocity of the deep HI 21cm absorption (large filled triangle at -30 km s^{-1} from Morganti et al. 2001), the shifts of the [O II] and [O III] lines from Tadhunter et al. (2001) (large filled diamonds at -5 km s^{-1} and -550 km s^{-1} respectively), the 2 components of the [O III] model in the nucleus (large open and filled squares) and the broad components of Pa α listed in Table 2 (open 8-pointed stars). For PA-5, positive positions are to the N and negative positions are to the S. For PA25, positive positions are to the SSW and negative positions are to the NNE.

3.2.3 Ionization and physical conditions

Given the success of the two component emission line profile described in the last section, it is possible to investigate the ionization, reddening and physical conditions for the narrow and intermediate emission line components detected in the nuclear aperture separately. The emission line ratios for both of the components derived from the double Gaussian profile fits are shown in the second column of Table 3. In addition to the nuclear aperture, Table 3 also shows emission line ratios determined for a 1.3×1.3 arcsecond aperture centred 1.6 arcseconds to the east of the nucleus along PA75

(AP1: VLT spectrum). In the case of this extended aperture, which is affected by seeing disk spillover of the nuclear emission line flux, it proved necessary to use a double Gaussian fit to model the lines accurately – one component representing the nuclear intermediate component and the other the extended narrow component. However, despite the contamination by the nuclear flux in this aperture, the narrow component has a much better contrast relative to the intermediate component than in the nuclear aperture; in Table 3 we only list the narrow line ratios for this aperture.

It is immediately clear from Table 3 that the




Figure 6. Double Gaussian model for the [O III] $\lambda\lambda 4959, 5007$ doublet in the nucleus, forcing the redshift of the narrow component to be consistent with the redshift of the narrow component in the low ionisation lines. The composite model is overplotted (bold line) on the extracted spectrum (faint line). All of the 4 components (2 for each line) are also plotted: the dotted lines trace the narrow components and the dashed lines trace the broad components.

blueshifted, intermediate component detected in the nucleus is characterised by a high ionization state, with relatively strong [NeV], [NeIII], HeII, [OIII] and [FeVII] lines, and relatively weak [OII], [OI] and [SII] lines.

In contrast, the ionization state of the narrow component detected in both the nuclear and extended apertures is relatively low, with [OII], [OI], [NII] and [SII] all strong relative to the higher ionization [NeV], [NeIII] and HeII and [OIII] lines.

On the basis of the $H\alpha/H\beta$ ratio alone there is no strong evidence for a high degree of reddening affecting the narrow line region in this object, in contrast to the situation in PKS1345+12 – an object that also shows broad, blueshifted Balmer lines (Holt et al. 2003). However, it is difficult to be entirely confident of the accuracy of the $H\alpha$ flux because of the complexity of the 7 Gaussian model required to fit the $H\alpha$ + [NII] blend. Moreover, the higher order Balmer lines cannot be used to check the reddening estimated from $H\alpha/H\beta$ because of their weakness relative to the continuum and underlying Balmer absorption line features. Given these uncertainties we have made no attempt to correct the emission line ratios for reddening.

Similarly, it has not been possible to estimate accurate electron temperatures for the narrow line gas using the [OIII](5007+4959)/4363 line ratio, because the faint [OIII] $\lambda 4363$ line is highly sensitive to the accuracy of the continuum subtraction, particularly for the broader, blueshifted component that is also blended with $H\gamma$. Nonetheless it has been possible to estimate the electron density from the [SII](6731/6717) ratio for the narrow emission line component detected in the nuclear aperture and the extended aperture along PA75. Using a model for the [SII] $\lambda\lambda 6717, 6731$ blend that includes both an intermediate blueshifted component and a narrow component for each line in the doublet, and assuming an electron temperature of 10,000K for the region emitting the [SII] lines, we

estimate an electron density $430 \pm 50 \text{ cm}^{-3}$ for the narrow component in the nuclear aperture. Unfortunately, because of the relative weakness of the intermediate blueshifted component to the lines, the electron density in the gas emitting the blueshifted component is not well-constrained by these observations. We find that, although 4 Gaussian models with the doublet ratios left unconstrained tend to the high density limit for the intermediate component, models with the ratio of the intermediate component fixed at the low density limit also provide adequate fits to the blend. On the other hand, the ratios determined for the narrow components are not sensitive to the details of the model used for the intermediate components.

In order to investigate the ionization mechanism(s) for the various kinematic components, the emission line ratios are compared with various ionization models and the results for other radio galaxies in the four diagnostic diagrams presented in Figure 7. Four sets of models are plotted: single slab optically thick power law photoionization models calculated using the MAPPINGS code; mixed medium photoionization models taken from Binette et al. (1996); pure shock cooling zone models and shock plus precursor models from Dopita & Sutherland (1995, 1996).

These diagnostic diagrams further emphasise the high ionization character of the gas emitting the intermediate, blueshifted component. In all diagnostic diagrams except that involving [NII], the line ratios for this component can only be fitted with power-law AGN photoionization models that have a high ionization parameter ($0.05 < U < 0.1$), and the line ratios are most consistent with models that have a power-law spectral index of -1.5. Although the particular mixed medium photoionization models plotted in the diagrams do not provide a good fit to the line ratios of the intermediate component, it is possible that, by changing the parameters of such models (e.g. ionization parameter, column depth of matter bounded component), an acceptable fit could be obtained. It is also notable that none of the shock or shock+precursor can provide an adequate fit to the line ratios.

The intermediate component has line ratios that place it beyond the high ionization end of the locus of narrow line radio galaxies, overlapping with the range of ionization parameter measured for the narrow, blueshifted absorption line systems detected against the UV and X-ray continua of Seyfert 1 galaxies and quasars (Crenshaw et al. 2003). Indeed we would expect such absorption line systems to be detected in PKS1549-79 for lines of sight to the AGN that intercept the high ionization gas. The only line ratio that does not fit in with this general trend is [NII]/ $H\alpha$, which is unusually large considering the high ionization state revealed by the other lines. This may reflect the fact that the systematic uncertainties for this ratio are larger than for the others because of the complexity of the $H\alpha$ + [NII] blend. Moreover, the models plotted in Figure 7 assume Solar elemental abundance ratios, whereas the abundance of nitrogen relative to other elements such as oxygen may be significantly larger than the Solar value – the favoured explanation for the unusually strong [NII] in, for example, Cygnus A (Tadhunter, Metz & Robinson 1994).

The diagnostic diagrams also show that the line ratios of the narrow emission line components (all apertures) fall at the lower ionization end of the ratios measured for the




Figure 7. Emission line diagnostic diagrams for PKS 1549-79. *Upper left:* $[\text{OIII}]/\text{H}\beta$ vs. $[\text{OII}]/[\text{OIII}]$. *Upper right:* $[\text{OIII}]/\text{H}\beta$ vs. $[\text{NII}]/\text{H}\alpha$. *Bottom left:* $[\text{OII}]/[\text{OIII}]$ vs. $[\text{NeV}]/[\text{NeIII}]$. *Bottom right:* $[\text{OIII}]/[\text{OII}]$ vs. $[\text{OI}]/[\text{OIII}]$. In these diagrams, the pure shock models of Dopita & Sutherland (1996) for a range of magnetic parameter ($B/\sqrt{n} = 0, 1, 2 \mu\text{G cm}^{3/2}$) with shock velocities $150 \leq v_{\text{shock}} \leq 500 \text{ km s}^{-1}$ are represented by dotted lines, while the shock plus precursor models are represented by dashed lines. Solid lines represent optically thick power law ($F_{\nu} \propto \nu^{\alpha}$) photoionization models calculated using the MAPPINGS code for three power-law spectral indices ($\alpha = -1.0, -1.5, -2.0$) for a sequence in the ionization parameter ($2.5 \times 10^{-4} \leq U \leq 10^{-1}$) increasing from right to left. Most of the photoionization models plotted in these figures were calculated for low electron densities ($N = 10^2 \text{ cm}^{-3}$), however the dot dashed lines shows a sequence of models calculated at a single ionization parameter ($U = 5 \times 10^{-3}$) for a range of densities ($10^2 < N < 10^6 \text{ cm}^{-3}$). The bold line corresponds to mixed-medium models from Binette et al. (1996) that include both matter-bounded and ionization-bounded clouds for a range of ratios of the solid angles covered by the matter bounded and ionization bounded clouds ($10^{-2} \leq A_{\text{M}/\text{I}} \leq 10$) increasing from right to left. The nuclear intermediate component is represented by the large solid square, while the narrow components for the nuclear and extended apertures are represented by large solid and open triangles respectively. For comparison, the small filled triangles represent data for the nuclear narrow line regions of radio galaxies (data from: Wills et al. 2002, Storch-Bergmann et al. 1996, Tadhunter 1987, Grandi & Osterbrock 1978, Costero & Osterbrock 1977, Cohen & Osterbrock 1977, Cohen & Osterbrock 1981, Dickson, 1997, Solorzano-Inarrea 2001, Villar-Martin et al. 1998, Clark 1996, Robinson et al. 2000), small filled squares represent extended emission line regions (data from: Tadhunter et al. 1994, Storch-Bergmann et al. 1996, Morganti et al. 1991, Robinson et al. 2000), and open squares represent EELRs with strong jet-cloud interactions (data from: Villar-Martin et al. 1998, 1999, Clark 1996, Clarke et al. 1998, Solorzano-Innarea 2001). On addition, we present the line ratios for various kinematic components detected in the spatially resolved NLR of Cygnus A by Taylor, Tadhunter & Robinson (2003), with the nuclear broad and narrow components plotted as small open triangles, and the extended narrow components plotted as open diamonds.




Figure 8. Near-IR (K-band) spectrum of PKS1549-79 taken using the IRIS2 spectrograph on the AAT. The main spectral features are marked. At near-IR wavelengths the spectrum is dominated by a quasar component.

narrow line regions of radio galaxies in general, consistent with photoionization at moderate ionization parameter, or shock+precursor ionization models. Despite the evidence for young stellar populations in this source, and the possibility of on-going star formation, the strengths of the [OI] and [NII] emission lines relative to H α preclude HII region-like photoionization by normal OB stars.

3.3 Near-IR spectroscopy

Our new deep K-band spectrum of the nuclear regions taken with the ATT/IRIS2 is shown in Figure 8. This represents a substantial improvement over the previous K-band spectrum published in Bellamy et al. (2003), which was affected by a faulty grism that resulted in a splitting of all spectral features. As well as the strong Pa α emission line, the spectrum shows a feature near $2.26\mu\text{m}$ that is likely to represent a blend of H $2\nu=1-0\text{S}(3)$ and [SiVI] $\lambda 1.963\mu\text{m}$, an H $2\nu=1-0\text{S}(2)$ line and (less certainly) a broad feature close to the expected wavelength of redshifted Br δ .

The Pa α feature can be modelled as a combination of two Gaussians of width $1220\pm 90\text{ km s}^{-1}$ and $3540\pm 320\text{ km s}^{-1}$ (FWHM) with fluxes of $(1.10\pm 0.16)\times 10^{-14}\text{ erg cm}^{-2}\text{ s}^{-1}$ and $(1.92\pm 0.15)\times 10^{-14}\text{ erg cm}^{-2}\text{ s}^{-1}$ respectively; redshifts for these components are given in Table 2. Similar results are obtained from fitting our NTT/SOFI spectrum. Although the narrower of these two broad components has a width similar to that of the intermediate component detected in the [OIII] lines, its redshift is close to that of the narrow emission line components (see Table 2), therefore it cannot be emitted by the same gaseous system that emits the intermediate [OIII] line component. Given this, and the fact that the broader Pa α component has a width that falls outside the normal range for the narrow line gas associated with AGN, it is likely that the Pa α line is dominated by genuine broad line region (BLR) emission. For comparison, a single Gaussian fit to the Pa α gives a linewidth: FWHM = $1940\pm 50\text{ km s}^{-1}$.

The new spectra also allow us to resolve the uncertainty in the identification of the feature at $\sim 2.26\mu\text{m}$ in the observed spectrum (see Figure 8). In terms of its red-

Emission line	Comp.	Nuclear flux	Extended flux
(a)	(b)	(c)	(d)
[Ne V] 3425.9	n	-	< 20
	i	75 ± 8	-
[O II] 3727.64†	n	330 ± 30	420 ± 50
	i	38 ± 8	-
[Ne III] 3868.8	n	50 ± 8	80 ± 10
	i	110 ± 8	-
[Ne III] 3968.4	n	18 ± 3	26 ± 4
	i	38 ± 4	-
H γ 4339.95	n	33 ± 4	-
	i	27 ± 3	-
[O III] 4363.2	n	8 ± 3	< 22
	i	< 8	-
He II 4685.75	n	11 ± 4	20 ± 10
	i	52 ± 7	-
H β 4860.75	n	100	100
	i	100	-
[O III] 4958.9	n	140 ± 10	150 ± 30
	i	580 ± 45	-
[O III] 5006.9	n	410 ± 50	450 ± 80
	i	1700 ± 150	-
N I 5199.1	n	51 ± 6	-
	i	< 10	-
He I 5875.2	n	64 ± 8	-
	i	< 10	-
[Fe VII] 6086.0	n	< 10	-
	i	42 ± 5	-
[O I] 6300.3	n	160 ± 20	80 ± 10
	i	57 ± 10	-
[O I] 6363.8	n	53 ± 6	28 ± 4
	i	15 ± 4	-
[N II] 6548.09	n	420 ± 30	180 ± 20
	i	190 ± 20	-
H α 6562.9	n	400 ± 50	500 ± 70
	i	350 ± 40	-
[N II] 6583.36	n	1230 ± 100	560 ± 70
	i	570 ± 100	-
[S II] 6716.4	n	290 ± 15	120 ± 20
	i	20 ± 4	-
[S II] 6730.8	n	270 ± 15	140 ± 20
	i	47 ± 10	-
F(H β)	n	2.4×10^{-16}	0.9×10^{-16}
(erg cm $^{-2}$ s $^{-1}$)	i	5.4×10^{-16}	-

Table 3. Data for all optical emission lines detected in the nuclear and extended apertures of PKS 1549-79. Columns are: (a) emission line; (b) emission line component where n and i are the narrow and intermediate components; (c) emission line fluxes measured for a 1.3×1.3 arcsecond aperture centred on the nucleus, extracted from the PA75 VLT data (fluxes expressed relative to H β); (d) emission line fluxes measured for a 1.3×1.3 arcsecond extended aperture centred 1.6 arcseconds east of the nucleus, extracted from the PA75 VLT data.

† weighted mean of [O II] doublet at the low density limit.




Figure 9. Fit to the shape of the near-IR H+K spectrum of PKS1549-79 taken on the NTT, showing the extrapolation to optical wavelengths for various assumptions about the power-law shape of the optical-IR continuum. Also shown are the VLT optical spectrum (upper curve at short wavelengths), and the VLT spectrum following subtraction of the best fitting stellar continuum model (lower curve at short wavelengths). The upper and lower dashed lines represent models with the minimum and maximum reddening respectively, while the solid line represents the average reddening case (see Table 4). The arrow in the lower left corner of the plot represents the upper limit on the AGN continuum flux at 5900Å derived from HST imaging observations.

shift this feature could either be identified with blueshifted [SiVI] λ 1.963 μ m or with $H_2\nu=1-0S(3)$ at the redshift of the nuclear narrow component. However, a single Gaussian fit to this feature gives a width of $970 \pm 80 \text{ km s}^{-1}$, which is significantly broader than the nuclear narrow component, but significantly narrower than the intermediate component, suggesting that the feature is composite. Therefore we have attempted a double Gaussian fit to the feature, with one Gaussian component representing $H_2\nu=1-0S(3)$ with the same width and redshift as the nuclear narrow component, and the other representing [SiVI] λ 1.963 μ m with the same width and redshift as the nuclear intermediate component. This double Gaussian model provides a good fit to the data with fluxes of $(4.5 \pm 1.5) \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$ and $(3.1 \pm 0.3) \times 10^{-15} \text{ erg cm}^{-2}$ for the narrow $H_2\nu=1-0S(3)$ and intermediate [SiVI] λ 1.963 μ m components respectively. The detection of $H_2\nu=1-0S(3)$ is consistent with the presence of $H_2\nu=1-0S(2)$ in the spectrum at longer wavelengths; the estimated luminosity for the $H_2\nu=1-0S(3)$ line ($2.3 \times 10^{40} \text{ erg s}^{-1}$) is less than estimated by Bellamy et al. (2003) who assumed that all the $\sim 2.26\mu\text{m}$ feature is $H_2\nu=1-0S(3)$. It is notable that the detection of the blueshifted [SiVI] λ 1.963 μ m line is consistent with the high ionization character of the

intermediate emission line component deduced from the optical observations (see section 3.2).

Along with information about the emission line systems, the new spectra provide key information about the quasar continuum emission. In particular, the new NTT spectrum has a broader spectral coverage than the AAT/IRIS spectra and allows more stringent limits to be placed on the reddening to the AGN. To estimate the AGN reddening we follow the same technique as Bellamy et al. (2003): fitting the near-IR continuum spectrum assuming a range of intrinsic power-law optical-infrared spectral shapes for quasars taken from Simpson & Rawlings (2000). The results are presented in Table 4 and Figure 9.

On the basis of this modelling alone it is clear that the V-band absorption and absolute magnitude of the AGN must fall in the range $4.9 < A_v(\text{intrinsic}) < 13.2$ and $-27.56 < M_v < -22.64$ respectively. However, the recent HST/ACS imaging of PKS1549-79 (Batcheldor et al. 2006 in preparation) allows us to place a stringent upper limit on the flux of the quasar core at 5900Å(rest) of $F_\lambda < 1.0 \times 10^{-17} \text{ erg cm}^{-2} \text{ Å}^{-1}$. Using this limit it is clear from Figure 9 that the intrinsic absorption and luminosity of the quasar must lie at the upper end of the range predicted on the basis of

	Minimum	Average	Maximum
$\alpha(\text{intrinsic})$	1.62	0.90	-0.67
E(B-V)	1.52	2.31	4.04
A_v	4.94	7.52	13.19
M_v	-22.64	-24.18	-27.56
M_k	-26.97	-27.43	-28.44

Table 4. Reddening and absolute magnitude estimates for the quasar nucleus in PKS1549-79. The second, third and fourth columns give the minimum, average and maximum reddening corresponding to the range of optical/UV power-law slopes deduced by Simpson & Rawlings (2000). The second row gives the values of the assumed spectral index for the intrinsic optical-IR spectral shape (convention: $F_\nu \propto \nu^{-\alpha}$). The final four rows give estimates for the E(B-V) reddening, A_v extinction, and the estimated absolute magnitude of the quasar nucleus in both the V and the K bands. In modelling the continuum shape we have assumed the extinction law of Rieke & Lebofsky (1985).

the near-IR continuum modelling; we find that the optical flux limit implies $A_v > 6.4$, $M_V < -23.5$ and $M_K < -27.3$. This places the AGN in PKS1549-79 firmly in the luminosity range of quasars ($M_v < -23.0$). Note also that such high values of nuclear extinction and luminosity are consistent with the residual spectrum obtained by subtracting the best fitting stellar continuum model from our VLT spectrum (see Figure 4).

3.4 Radio imaging and spectroscopy

3.4.1 VLBI continuum imaging

Figure 10 shows the VLBI images made with SHEVE at 2.3 and 8.4GHz. The radio source is characterised by a compact core to the western extreme of the source, which shows a jet-like extension in PA64, and a highly collimated jet structure to the east aligned along PA120. The overall extent of the source in the 2.3GHz image is 180mas (430 pc), and there is no strong evidence for emission from more extended structures; >95% of the total (single dish) flux is contained within the structures visible in the VLBI images. Relative to the western core the jet appears significantly fainter in the 8.4GHz image (core/jet ratio $\sim 10:1$) than it does in the 2.3GHz image (core/jet ratio $\sim 2:1$), but the western core has a similar brightness in both images. The non-simultaneity of the 2.3GHz and 8.4GHz VLBI data, and the resolution differences between the two frequencies, preclude accurate determination of the power-law spectral indices of the jet and core components. However, the data are consistent with a steep spectrum for the jet ($\alpha \sim -1.2$; $F_\nu \propto \nu^{+\alpha}$) and a much flatter spectrum ($\alpha \sim 0$) for the core. Note that the total flux recovered in our VLBI observations at 8.4 GHz (3.6 Jy — data taken in 1991) is significantly larger than the 8.4 GHz flux measured more recently by Drake et al. (2004b: 2.4 Jy — data taken between 1996 and 1998). This suggests that the core is variable at high frequencies.

Overall, the radio structure is similar to the core-jet structures detected in some other radio-loud quasars such

as 3C273. Such structures are generally interpreted in terms of radio emission from a two sided relativistic jet structure, with the axis of jets pointing close to our line of sight. In this case the radio emission from the jet on the near side is enhanced because the radio emission is beamed towards the observer, whereas the emission from the jet on the far side is dimmed because the radiation is beamed away from the observer. The significant curvature in the jet is also consistent with the beaming hypothesis, since a small degree of intrinsic jet curvature will be exaggerated if the jet is pointing close to the line of sight. If this beaming/orientation explanation for the one-sided jet is correct, then the total radio emission from the source will be boosted relative to the situation in which the jet is close to the plane of the sky. Therefore, although the total radio power of PKS1549-79 uncorrected for beaming ($P_{2.3\text{GHz}} = 5.4 \times 10^{26} \text{ W Hz}^{-1}$) would lead to its classification as a powerful radio galaxy that would normally appear as an FR II radio source if orientated close to the plane of the sky, the true (unbeamed) radio power of the source could be much lower. If we assume that the radio flux not recovered in the VLBI maps (i.e. 5% of the total) represents an unbeamed radio component, this leads to a lower limit on the total unbeamed radio power of $P_{2.3\text{GHz}} = 2.7 \times 10^{25} \text{ W Hz}^{-1}$ — just above the break in the radio luminosity function (Auriemma et al. 1977) and in the transition region between FRI and FR II sources (Fanaroff & Riley 1974).

Assuming that the one sided nature of the jet is a consequence of beaming effects, we can use the lower limit on the jet to counter-jet ratio (R_{jc}) derived from the radio maps to estimate the inclination of the jet to the line of sight. The 21cm LBA continuum map (see Figure 11) is the most sensitive for the detected of extended jet features close to the nucleus. The peak flux in the jet detected in this map is 2.3 Jy beam $^{-1}$, whereas the 3σ detection limit for the counter jet is 48 mJy beam $^{-1}$. Therefore a lower limit on the jet to counter-jet ratio is $R_{jc} \geq 48$. The jet to counter-jet ratio is related to the bulk Lorentz factor in the jet (γ) and the inclination of the jet to the line of sight (i) by:

$$\cos(i) = \left(\frac{\gamma^2}{\gamma^2 - 1} \right)^{1/2} \left(\frac{R_{jc}^{1/(2-\alpha)} - 1}{R_{jc}^{1/(2-\alpha)} + 1} \right) \quad (1)$$

where α is the spectral index of the jet continuum emission. For $\alpha = -1$ and $R_{jc} \geq 48$ we estimate an upper limit on the inclination of $i < 55$ degrees, assuming that the Lorentz factor for the bulk motion of the jet falls within the typical range estimated for extragalactic radio jets ($5 < \gamma < 10$).

An alternative possibility is that the one-sided jet is a consequence of a stronger interaction between the radio jet and the interstellar medium on the east side of the nucleus that leads to local re-acceleration of the electrons and enhanced radio emission; a strong jet-cloud interaction would also explain the curvature in the jet. However, this scenario would not account for the relative strength of the flat spectrum core and the fact that the asymmetric character of jet extends into the near-jet close to the core, as is clear from the higher resolution 2.3GHz map (Figure 10 middle).




Figure 10. High resolution VLBI radio maps of PKS1549-79. The upper map is for a frequency of 2.3GHz excluding the Hartebeesthoek antenna, the middle map is also for a frequency of 2.3 GHz but including the Hartebeesthoek antenna, while the lower map is for an observation frequency of 8.4 GHz (contours: -10,10,20,40,80,160,320,640,1280 mJy/Beam).

3.4.2 HI 21cm kinematics

Given the core-jet character of the radio continuum structure, which suggested that the jet is pointing close to our line of sight, it was a surprise to detect HI absorption in our ATCA observations (Morganti et al. 2001). On the basis of the simplest versions of the unified schemes one would normally expect a relatively unobscured view of the quasar nucleus from the direction of the radio jet (Barthel 1989). The absorption in the spatially integrated radio emission has a FWHM $\sim 80 \text{ km s}^{-1}$ and the peak optical depth is about 2%. This leads to a column density of $(4.0 \pm 0.5) \times 10^{18} T_{\text{spin}} \text{ cm}^{-2}$, where T_{spin} is the HI spin temperature. A possible interpretation described in Tadhunter et al. (2001) is that the HI absorption is associated with a dusty cocoon around the nucleus that also emits the narrow emission line component, and is responsible for the obscuration of the quasar nucleus.

The new LBA data provide information about the spatial structure of the HI absorption. Figure 11 shows the pro-

files of the HI absorption measured at the locations of both the core and the jet structures. Consistent with the idea of a foreground absorbing screen proposed by Morganti et al. (2001) and Tadhunter et al. (2001), the HI absorption was detected against both of the major radio components with similar redshift, linewidth and optical depth. Therefore the diameter of the HI absorbing region is > 120 milliarcseconds ($> 286 \text{ pc}$). Note that the narrow HI absorption is significantly blueshifted (by $\sim 75 \text{ km s}^{-1}$) relative to the narrow optical emission lines (see Figure 4 and Table 2), but this blueshift is an order of magnitude smaller than that measured in the [OIII] lines, and might be attributed to gravitational motions if the absorbing cloud has not yet settled into the circumnuclear disk.

In Figure 12 the HI absorption profile from the ATCA observations (from Morganti et al. 2001) is superimposed on the integrated absorption profile obtained from the new LBA observations. This shows that most of the HI absorption is recovered in the high spatial resolution LBA observations, and that the HI kinematics in the LBA and ATCA are consistent.

4 DISCUSSION

4.1 Evolutionary status and the triggering of the activity

The ready detection of multiple tidal tails at a relatively high surface brightness level is entirely consistent with the idea that the activity in PKS1549-79 has been triggered in a major merger between galaxies, at least one of which was gas rich and had a cold disk component prior to the merger. Indeed, the tail structure observed in PKS1549-79, with two near-parallel tails pointing to the south and a jet-like structure pointing to the north bears marked similarity with the structures produced in numerical simulations by Barnes & Hernquist (1996) of the merger between two disk galaxies, observed from the orbital plane of the merging galaxies. In such a merger, it will take of order 500 Myr for the nuclei of the two galaxies to finally merge together. Therefore it is likely that that we are observing the system a considerable period after the start of the encounter.

The young stellar populations (YSP) provide useful complementary information about the merger. The presence of a reddened YSP in the nuclear regions of this object is consistent with the idea that the merger also triggered a starburst. Given that the YSP has an estimated age of 50 – 250 Myr at the observation epoch, it is likely that the starburst was triggered a considerable period after the start of the merger sequence. This in turn suggests that one of the galaxies had a significant bulge that hindered the formation of a bar that would otherwise lead to the major merger-induced star formation occurring early in the merger (Mihos & Hearnquist 1996). Notwithstanding the uncertainties about the heating mechanism for the dust emitting the far-IR radiation, the substantial far-IR excess in this source suggests the presence of substantial ongoing star formation in the nuclear regions that is simultaneous with the currently observed quasar and jet activity.

It is also interesting to consider at what stage the radio source was triggered in the merger sequence. Tadhunter




Figure 11. *Middle:* 1.64 GHz LBA continuum image of PKS1549-79 obtained from the line-free channels. *Right:* integrated profile of the HI absorption in the western core source. *Left:* integrated profile of the HI absorption for the eastern jet source.

et al. (2001) have argued that, since the extreme forbidden emission line widths measured in PKS1549-79 are only detected in radio galaxies with young, compact radio sources, it is likely that we are observing the radio source at a relatively early stage after the triggering of the activity ($< 10^6$ yr). If this is correct, it is plausible that there was a significant delay (> 40 Myr) between the starburst associated with the YSP detected at optical wavelengths and the triggering of the radio jet activity. However, the evidence for such a time delay is weaker in PKS1549-79 than for some other radio sources in which the time lag between the the major episode of merger-induced star formation and the triggering of the currently observed radio source is estimated as 0.3 – 2.0 Gyr (Tadhunter et al. 2005, Emonts et al. 2005, Johnston et al. 2005).

4.2 The nature of the AGN

Our new data provide key information about the nature of the AGN in PKS1549-79. Most notably, the relatively narrow Pa α detected in the quasar component of PKS1549-79 (FWHM = 1940 ± 50 km s $^{-1}$ from a single Gaussian fit) is consistent with the classification of this source as a narrow line Seyfert 1 galaxy (NLS1: FWHM < 2000 km s $^{-1}$: Osterbrock & Pogge 1985, Osterbrock 1987) or more accurately, given its luminosity, a narrow line quasar. Interestingly, the extreme blueshift of the [OIII] line relative to the galaxy rest frame in this object is also similar to those detected in the broad [OIII] emission line components of many NLS1 galaxies and similar objects (Veron-Cetty & Veron 2001, Zamanov et al. 2002, Boroson 2005, Aoki, Kawaguchi & Ohta 2005).

NLS1 are often interpreted as AGN that are accreting at a relatively high Eddington ratio ($L_{bol}/L_{edd} \sim 1$), based on the small virial black hole mass estimates deduced from their relatively narrow permitted line widths (Pounds, Done & Osborne 1995, Mathur 2000, Mathur & Grupe 2005). It has also been suggested that the NLS1 fall significantly below the correlations between black hole mass and host galaxy




Figure 12. Profile of the HI absorption in PKS 1549-79. The solid line represents the ATCA data of Morganti et al. (2001), while the dotted lines represent the integrated HI profile measured from the new LBA data. Most of the absorbed flux is recovered in the higher resolution LBA observations.

bulge properties (Grupe & Mathur 2004). However, it is important to add the caveats that (a) the small permitted line widths of NLS1 may reflect orientation effects (e.g. BLR disk viewed face on) rather than low black hole masses; and (b) the bulge properties for NLS1 in the Grupe & Mathur (2004) study were deduced from [OIII] linewidths, which are likely to be affected by non-gravitational (radial) motions suggested by the large [OIII] blueshifts detected in some sources.

It is interesting to consider whether PKS1549-79 is also a high accretion rate object with a large Eddington ratio. Its black hole mass can be estimated using two independent methods. First, a virial mass estimate can be obtained using the Pa α line width and luminosity. Assuming the minimum extinction for Pa α ($A_k = 0.74$ magnitudes),

and a $\text{Pa}\alpha/\text{H}\beta$ ratio set at the case B recombination value ($\text{Pa}\alpha/\text{H}\beta = 0.332$, see Osterbrock 1989), we deduce an intrinsic quasar $\text{H}\beta$ luminosity of $L(\text{H}\beta) = 1.0 \times 10^{43} \text{ erg s}^{-1}$ for our adopted cosmology. Assuming that $\text{FWHM}(\text{H}\beta) = \text{FWHM}(\text{Pa}\alpha) = 1940 \text{ km s}^{-1}$ we then use the relationship between $\text{FWHM}(\text{H}\beta)$, $L(\text{H}\beta)$ and M_{bh} published by Greene & Ho (2005: their eqn 7) to estimate a virial black hole mass of $M_{bh} = 5.3 \times 10^7 M_\odot$. A second black hole mass estimate can be obtained from the correlation between R-band bulge luminosity (M_R) and black hole mass published by McClure and Dunlop (2002) along with the published R-band absolute magnitude for PKS1549-79 (Drake et al. 2004a: $M_R = -21.9$ for our cosmology)*. This leads to $M_{bh} = 2.6 \times 10^8 M_\odot$ — almost an order of magnitude larger than the virial estimate of black hole mass. However, the R-band luminosity estimate can be regarded as an upper limit because the YSP in the bulge and tail features of PKS1549-79 along with any AGN contamination are likely to artificially boost the R-band luminosity; the only way in which the R-band magnitude can substantially underestimate the black hole mass is if the host galaxy as a whole is subject to substantial intrinsic extinction. In contrast, the virial estimate is likely to represent a lower limit on the black hole mass because of the effects of orientation on the $\text{Pa}\alpha$ line width, especially given that the radio observations suggest that radio jet is pointing close to the line of sight, with the putative BLR disk observed close to face-on. The true mass of the black hole in PKS1549-79 is therefore likely to fall between the virial and bulge luminosity estimates.

The other element required to determine the Eddington ratio is the bolometric luminosity (L_{bol}) of the quasar. L_{bol} can be estimated using the quasar continuum modelling results along with information on generic quasar spectral energy distributions (SEDs) presented in Elvis et al. (1994). The range of optical SEDs that are consistent with our optical/IR spectroscopy and imaging have monochromatic V-band luminosities in the range $1.2 \times 10^{41} < L_V < 4.8 \times 10^{42} \text{ erg s}^{-1} \text{ \AA}^{-1}$ or $6.6 \times 10^{44} < \lambda L_\lambda < 2.6 \times 10^{46} \text{ erg s}^{-1}$ ($\lambda = 5500 \text{ \AA}$) for the quasar in PKS1549-79. The V-band bolometric correction factor listed in Elvis et al. (1994) is: $L_{bol}/(\lambda L_V) = 14.2$. Therefore, we estimate that the bolometric luminosity of PKS1549-79 falls in the range $9.0 \times 10^{46} < L_{bol} < 3.8 \times 10^{47} \text{ erg s}^{-1}$. Note that, although the radio jet in this source is pointing close to our line of sight, there is no evidence that the optical/IR continuum is substantially affected by contamination by a beamed synchrotron source: not only does the spectral energy distribution of PKS1549-79 show a steep decline between radio and sub-mm wavelengths (compare Drake et al. 2004b with Beasley et al. 1997), but both the $\text{H}\beta$ equivalent width and the L_{bol}/L_{ir} ratio fall well within the range measured for un-beamed quasars.

These estimates of L_{bol} lead to Eddington ratios in the range $0.3 < L_{bol}/L_{edd} < 11$ for our upper estimate of the

black hole mass ($M_{bh} = 2.6 \times 10^8 M_\odot$) and in the range $1.3 < L_{bol}/L_{edd} < 57$ for our lower estimate of the black hole mass ($M_{bh} = 5.3 \times 10^7 M_\odot$). The measurement of such high Eddington ratios reinforces the link between PKS1549-79 and NLS1 galaxies.

Although we cannot rule out the idea that PKS1549-79 is a significant outlier relative to the statistical correlations we have used to derive both black hole mass and bolometric luminosity estimates, our results are at least consistent with accretion at a high Eddington ratio in this object. On this evidence, accretion at high Eddington ratio does not prevent the formation of powerful, relativistic jets. For comparison, the Eddington ratio measured for Cygnus A — the only quasar with a direct, dynamical black hole mass estimate — is $(1.6 - 6.0) \times 10^{-2}$ (Tadhunter et al. 2003). Similarly, Dunlop et al. (2003) derive relatively small Eddington ratios (typically $L_{bol}/L_{edd} < 0.1$) based on R-band black hole mass estimates of several radio-loud and radio-quiet quasars. On the other hand, Punsly & Tingay (2005) have recently highlighted the high redshift quasar PKS0743-67 as another object that is accreting at high Eddington ratio but has powerful radio jets.

4.3 Characterising the warm outflow

Recent theoretical studies of galaxy evolution have invoked quasar-induced outflows to explain both the correlations between the masses of black holes and the galaxy bulge properties (e.g. Silk & Rees 1998, Fabian 1999, di Matteo et al. 2005), and the form of the galaxy luminosity function at the high luminosity end (e.g. Benson et al. 2003). Such feedback models require that a relatively large fraction of the available accretion power of the quasars is thermally coupled to the circumnuclear ISM ($\geq 10\%$: Fabian 1989, di Matteo et al. 2001).

PKS1549-79 appears to represent a situation close to that modelled in some of the recent simulations (e.g. di Matteo et al. 2005), in which the black hole is growing rapidly via merger-induced accretion following a merger between gas rich galaxies. We have also found — in the blueshifted [OIII] lines — clear evidence for an outflow in the central regions of this source. Therefore it is important to investigate whether the properties of the warm gas outflow in PKS1549-79 are consistent with the assumptions of the quasar feedback models.

We start by determining the mass outflow rate in the intermediate (blue shifted) component. For a steady state outflow the mass outflow rate is given by:

$$\dot{M} = N \epsilon v_{out} A m_p \quad (2)$$

where N is the electron density, ϵ the volume filling factor, v_{out} the outflow velocity, A the area of the outflow and m_p the mass of the proton. The filling factor is related to the $\text{H}\beta$ luminosity ($L(\text{H}\beta)$), density and emitting volume V as follows:

$$\epsilon = \frac{L(\text{H}\beta)}{\alpha_{H\beta}^{eff} h \nu_{H\beta} N V} \quad (3)$$

where $\alpha_{H\beta}^{eff}$ is the effective $\text{H}\beta$ recombination coefficient (see Osterbrock 1989), $\nu_{H\beta}$ the frequency of $\text{H}\beta$, and h the Planck constant. It follows that the mass outflow rate can be written as:

* Note that the R-band absolute magnitude derived by Drake et al. (2004a) is based in the integrated light measured within the outermost ellipse fitted to their R-band image (corresponding to a semi-major axis of 12.5 arcseconds or 30 kpc). Since this magnitude corresponds to the total light, and no disk/bulge separation has been attempted, it is likely to *overestimate* the luminosity of the bulge.

$$\dot{M} = \frac{L(H\beta)m_p v_{out} A}{\alpha_{H\beta}^{eff} h \nu_{H\beta} N V} \quad (4)$$

and for a spherical outflow this reduces to:

$$\dot{M} = \frac{3L(H\beta)m_p v_{out}}{N \alpha_{H\beta}^{eff} h \nu_{H\beta} r} \quad (5)$$

where r is the radius of the spherical volume. On the basis of our spectroscopy we estimate an $H\beta$ luminosity of $L(H\beta) = 2.8 \times 10^{40}$ erg s $^{-1}$ for the intermediate, blueshifted component and a rest frame outflow velocity of $v_{out} = 680$ km s $^{-1}$. If we assume case B recombination theory for an electron temperature $T = 10,000$ K and that the spherical outflow has the same radius as the radio source (180 milli-arcseconds or $r = 430$ pc for our cosmology) we find that the mass outflow rate in units of solar masses per year is given by:

$$\dot{M} = 9.1 \left(\frac{10^2 \text{ cm}^{-3}}{N} \right) M_{\odot} \text{ yr}^{-1}. \quad (6)$$

The largest uncertainty in this calculation is the electron density of the outflowing gas (N). Because it has not been possible to measure the electron density for the outflowing gas directly in the case of PKS1549-79, we assume that it falls in the range measured for the NLR of other radio galaxies. The typical electron densities measured for kpc-scale ionized gas in extended radio sources such as Cygnus A (Taylor, Tadhunter & Robinson 2003) and 3C321 (Robinson et al. 2000) are $N \sim 10^2$ cm $^{-3}$. However, the densities may be larger in the emission line regions associated with compact radio sources, and in the case of PKS1345+12 – a similar object that has broad, blueshifted [OIII] lines – a relatively high electron density ($N > 5 \times 10^3$ cm $^{-3}$) has been deduced for the blueshifted emission line components (Holt et al. 2003). Therefore it is reasonable to assume densities in the range $10^2 < N < 10^4$ cm $^{-3}$, leading to mass outflow rates $0.12 < \dot{M} < 12 M_{\odot} \text{ yr}^{-1}$. It is notable that this covers same range as the mass outflow rates deduced for the NLR of Seyfert galaxies (Veilleux, Cecil & Bland-Hawthorn 2005) and the high ionization absorption line systems detected at UV and X-ray wavelengths in some AGN (Crenshaw et al. 2003), but is somewhat less than deduced for the neutral outflows detected in extreme starburst sources such as ULIRGs (Rupke et al. 2005a,b). For the same range of densities we estimate that the total mass in the warm gas outflow falls in the range $1.9 \times 10^4 < M_{total} < 1.9 \times 10^6 M_{\odot}$ and the filling factor falls in the range $2.3 \times 10^{-3} < \epsilon < 2.3 \times 10^{-7}$.

From the point of view of gauging the likely impact of the warm gas outflow on the circumnuclear gas in the bulge of the host galaxy of PKS1549-79 it is also important to estimate the kinetic power of the outflow, including both the radial and turbulent components in the gas motion. Assuming that the relatively large linewidth of the outflowing gas reflects a turbulent motion that is present at all locations in the outflow region, the kinetic power is:

$$\dot{E} = 6.34 \times 10^{35} \frac{\dot{M}}{2} (v_{out}^2 + (FWHM)^2 / 5.55) \text{ erg s}^{-1} \quad (7)$$

where $FWHM$ is the full width at half maximum of the blueshifted [OIII] line in km s $^{-1}$, and \dot{M} is the mass outflow rate expressed in solar masses per year. For $v_{out} = 680$ km s $^{-1}$, $FWHM = 1282$ km s $^{-1}$ and the mass outflow rates

determined above, we find that the kinetic power for the outflowing gas falls in the range $2.2 \times 10^{40} < \dot{E} < 2.2 \times 10^{42}$ erg s $^{-1}$. Taking the Eddington luminosity derived from the larger black hole mass estimate (see section 4.2 above) as an estimate of the total power available from accretion, the kinetic power in the warm gas component represents only a small fraction ($7 \times 10^{-7} < \dot{E}/L_{edd} < 7 \times 10^{-5}$) of the available accretion power, in contrast to the quasar feedback models which require a much larger fraction of the accretion power of the black hole to be thermally coupled to the gas surrounding the black hole. It is also unlikely that the warm gas outflow by itself is capable of removing all the warm/cool gas from the central regions of the host galaxy. Assuming that PKS1549-79 has a total mass $M_{total} \sim 10^{11} M_{\odot}$, and a gas mass $M_{gas} \sim 10^{10} M_{\odot}$ contained within a radius of 5 kpc – conservative assumptions for ULIRGs which are likely to have a larger gas mass concentrated within a smaller radius – the gravitational binding energy of the gas is $E_{bind} \approx GM_{gas}M_{total}/R_{gas} \approx 2 \times 10^{58}$ erg. In comparison, the warm gas outflow will deposit only $\sim 10^{55} - 10^{57}$ erg into the surrounding ISM, assuming that it can persist in its current form for the typical 10^7 yr lifetime of an extragalactic radio source.

How do we explain the fact that the estimated kinematic power in the warm gas outflow in PKS1549-79 is at least three orders of magnitude less than might be expected on the basis of the quasar feedback models? Considering first the assumptions that have gone into the above estimates, the key parameters are the radius of the outflow, the electron density and the emission line luminosity. \dot{E} would be underestimated if the radius and/or density of the outflow have been overestimated, or the $H\beta$ luminosity has been underestimated. However, it is unlikely that the radius of the outflow region has been overestimated, because recent HST imaging observations of PKS1549-79 demonstrate that it has a similar scale to the radio source (Batcheldor et al. 2006); allowing for projection effects, it is more likely that the radius we have used is conservative and the outflow region is significantly *larger* than we have estimated. Similarly, given that warm gas has been accelerated to large velocities and other compact radio sources show strong evidence for a relatively high density NLR (Holt et al. 2003, Holt 2005), it is unlikely that the electron density in the outflow can be substantially less than the measured density in the narrow component (430 ± 50 cm $^{-3}$) and the lower limit ($N \geq 10^2$ cm $^{-3}$) we have assumed.

Given the fact that the quasar nucleus in PKS1549-79 is itself substantially obscured at optical wavelengths, it is less easy to dismiss the idea that a portion of the outflow region is highly extinguished at optical wavelengths so that the $H\beta$ luminosity is underestimated. However, it is unlikely that the $H\beta$ has been underestimated by more than a factor 50 because the blueshifted outflow component would then make a substantial contribution to the Pa α flux, which is not observed.

Most plausibly, the warm gas outflow we have detected in [OIII] represents only a small fraction of the total mass in the outflow, with the remaining mass locked in hotter or cooler phases of the ISM that are difficult to detect at optical wavelengths. Recently, Morganti et al. (2003, 2005a,b) have detected massive *neutral* outflows in several compact radio sources using observations of the 21cm line of neutral hy-

drogen. The mass outflow rates and energy fluxes contained within such neutral outflows are substantially larger than we have estimated for the warm gas outflow in PKS1549-79. It is notable that there is a hint of a blue wing to the HI 21cm line detected in PKS1549-79 (see Figures 11 and 12). If this can be confirmed with higher quality observations, it may be possible to make a more complete census of the warm and cool gas outflows in this object.

4.4 HI spin temperature

The exact value of the HI spin temperature (T_{spin}) is a major uncertainty in calculating the column density of neutral hydrogen using HI 21cm absorption line observations. Although T_{spin} is often assumed to have values consistent with those estimated in the Galaxy ($100 < T_{\text{spin}} < 1000$ K; Heiles & Kulkarni 1988), higher values may be appropriate in the central regions around AGN because of excitation of the upper level of the 21cm hyperfine transition by collisions, Ly α photons or 21cm continuum photons (see Bahcall & Ekers 1969 for a discussion). The latter may be particularly important in a case like PKS1549-79 because the radio emission is beamed towards the HI absorbing cloud along our line of sight. However, to our knowledge there have so far been no direct estimates of the HI 21cm spin temperature in the vicinity of luminous AGN.

PKS1549-79 provides an excellent opportunity to estimate the spin temperature directly because we have an independent estimate of the neutral column towards the quasar nucleus from the extinction estimates derived from our quasar continuum modelling. The SED modelling gives $2.0 < E(B-V) < 4.0$ for the intrinsic line of sight reddening towards the quasar nucleus. Assuming the standard Galactic dust to gas ratio ($E(B-V)/N_{\text{HI}} = 1.7 \times 10^{-22}$ mag cm $^{-2}$; Bohlin et al. 1978) this translates into neutral HI columns in the range $1.2 \times 10^{22} < N_{\text{HI}} < 2.4 \times 10^{22}$ cm $^{-2}$. On the basis of our HI 21cm absorption line measurements, the neutral HI column is related to the spin temperature by: $N_{\text{HI}} = (4.0 \pm 0.5) \times 10^{18} T_{\text{spin}}$ cm $^{-2}$. Therefore, by combining the estimates of the neutral HI column derived from both the HI 21cm and the quasar SED measurements we find that the electron spin temperature must fall in the range $3000 < T_{\text{spin}} < 6000$ K. These measurements provide clear evidence that the HI spin temperature in PKS1549-79 is higher than the typical Galactic value.

Implicit in our estimate of T_{spin} is the assumption that the column reddening the quasar is the same as that causing the HI 21cm absorption. Although the quasar nucleus can be considered point-like, while the radio emission is clearly spatially extended in our VLBI observations, our LBA observations show that the HI absorption covers the entire spatially-resolved radio source, including the flat spectrum core to the western jet of the source. Therefore it is entirely plausible that the HI 21cm and quasar reddening columns are associated with the same cloud. Another major assumption we have made is that the dust-to-gas ratio in the absorbing cloud has the Galactic value. This is called into question by recent observations that provide evidence for dust-to-gas ratios in the vicinity of AGN that are a factor ~ 10 or more *lower* than the Galactic value (Maiolino et al. 2001). However, if such low dust-to-gas ratios also hold in PKS1549-79, the resulting spin temperature would be even larger than

we have estimated above, further strengthening the evidence for large T_{spin} in this object. In view of this the lower limit we have derived for the spin temperature in PKS1549-79 ($T_{\text{spin}} > 3.0 \times 10^3$ K) is likely to be conservative.

5 CONCLUSIONS AND FURTHER WORK

The new observations presented in this paper confirm that PKS1549-79 represents a luminous quasar observed at an early stage of its evolution. The quasar and jet activity have been triggered in a major merger which has also triggered substantial star formation. Accreting at close to the Eddington rate, the black hole in this system is growing rapidly and simultaneously driving relativistic jets. The growth phase of the black hole is substantially obscured at optical wavelengths in this object, consistent with recent galaxy evolution models (Hopkins et al. 2005). However, although the optical emission lines provide evidence for a high ionization, warm outflow associated with the AGN, the warm outflow currently observed at optical wavelengths is not in itself sufficient to remove all the gas from the central regions of the galaxy.

PKS1549-79 is a key object because it links together several classes of active galaxies, including radio galaxies, quasars, ULIRGs, NLS1 and AGN with high ionization absorption line systems. Moreover it shows many properties in common with the recently discovered population of high redshift AGN detected at sub-mm wavelengths (Alexander et al. 2005). The common thread is that all these classes of object are associated with the triggering and feedback effects of major galaxy mergers. Although much attention is paid to the links between star formation and activity in objects at high redshift, this study of PKS1549-79 serves to emphasise that some galaxies are evolving rapidly in the local universe, and that it is possible to use such objects to study the coevolution of black holes and galaxy bulges in depth.

Several further observational studies of PKS1549-79 would enhance our understanding of the relationship(s) between AGN activity and galaxy evolution. They include the following.

- Near-IR measurements of the stellar bulge luminosity and velocity dispersion to improve estimates of the black hole mass.
- High quality optical spectroscopy observations of a wider range of emission line diagnostics to measure the electron density accurately, and thereby improve our determination of the properties of the warm gas outflow.
- Deep X-ray spectroscopy observations to refine estimates of the quasar bolometric luminosity and absorbing HI column. Such observations would also allow the links between PKS1549-79, NLS1 and sub-mm galaxies to be further investigated.
- Multi-epoch VLBI observations to search for superluminal motions in the jet, and accurately determine the inclination of the jet to the line of sight.
- Deeper HI 21cm measurements and high resolution optical observations of the NaID line to investigate the importance of neutral gas outflows.

Acknowledgments

Based on observations collected at the European Southern Observatory, Chile (ESO Programmes 69.B-0548(A), 71.B-0320(A) and 72.B-0792(A)) and the Anglo Australian Observatory, Siding Spring, Australia. KJI, JH and MB acknowledge financial support from PPARC. Edward King (CSIRO) provided original images made with the SHEVE network. We thank the anonymous referee for useful comments. The LBA is part of the Australia Telescope, funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO and the University of Tasmania.

REFERENCES

- Alexander, D. M., Smail, I., Bauer, F. E., Chapman, S. C., Blain, A. W., Brandt, W. N., Ivison, R. J., 2005, *Nat*, 434, 738
- Aoki, K., Kawaguchi, T., Ohta, K., 2005, *ApJ*, 618, 601
- Areteaga, I., Terlevich, E., Terlevich, R.J., Cotter, G., Diaz, A.I., 2001, *MNRAS*, 325, 636
- Auriemma, C., Perola, G.C., Ekers, R.D., Fanti, R., Lari, C., Jaffe, W.J., Ulrich, M.H., 1977, *A&A*, 57, 41
- Bahcall, J.N., Ekers, R.D., 1969, *ApJ*, 157, 1055
- Barnes, J.A., Hernquist, L., 1996, *ApJ*, 471, 115
- Barthel, P.D., 1989, *ApJ*, 336, 606
- Baum, S.A., Heckman, T., van Breugel, W., 1990, *ApJS*, 74, 389
- Beasley, A.J., Conway, J.E., Booth, R.S., Nyman, L.-A., Holdaway, M., 1997, *A&ASuppl*, 124, 469
- Bellamy, M.J., Tadhunter, C.N., Morganti, R., Wills, K.A., Holt, J., Taylor, M.D., Watson, C.A., 2003, *MNRAS*, 344, L80
- Benson, A.J., Bower, R.G., Frenk, C.S., Lacey, C.G., Baugh, C.M., Cole, S., 2003, *ApJ*, 599, 38
- Bessell, M.S., Castelli, F., Plez, B., 1998, *A&A*, 333, 231
- Binette, L., Wilson, A.S., Storchi-Bergmann, T., 1996, *A&A*, 312, 365
- Bohlin, R.C., Savage, B.D., Drake, J.F., 1978, *ApJ*, 224, 132
- Boroson, T., 2005, *ApJ*, 130, 381
- Bruzual, G., Charlot, S., 2003, *MNRAS*, 344, 1000
- Clark, B.G., 1973, *Proc.IEEE*, 61, 1242
- Clark, N., 1996, PhD thesis, University of Sheffield
- Clark, N.E., Axon, D.J., Tadhunter, C.N., Robinson, A., O'Brien, P., 2005, *ApJ*, 494, 546
- Costero, R., Osterbrock, D.E., 1977, *ApJ*, 211, 675
- Crenshaw, D.M., Kraemer, S.B., George, I.M., 2003, *ARA&A*, 2003
- Dickson, R., Tadhunter, C., Shaw, M., Clark, N. & Morganti, R., 1995, *MNRAS*, 273, L29
- Dickson, R.D., 1997, PhD thesis, University of Sheffield
- di Matteo, T., Springel, V., Hernquist, L., 2005, *Nat*, 433, 604
- Dopita, M., Sutherland, R.S., 1995, *ApJ*, 455, 468
- Dopita, M., Sutherland, R.S., 1996, *ApJS*, 102, 161
- Drake, C.L., McGregor, P.J., Dopita, M.A., 2004a, *AJ*, 128, 955
- Drake, C.L., Bicknell, G.V., McGregor, P.J., Dopita, M.A., 2004b, *ApJ*, 128, 969
- Dunlop, J.S., Peacock, J.A., 1990, *MNRAS*, 247, 19
- Dunlop, J.S., McClure, R.J., Kukula, M.J., Baum, S.A., O'Dea, C.P., Hughes, D.H., 2003, *MNRAS*, 340, 1095
- Emonts, B.H.C., Morganti, R., Tadhunter, C.N., Holt, J., Oosterloo, T.A., van der Hulst, J.M., Wills, K.A., 2005, *A&A*, submitted
- Elvis, M., Wilkes, B.J., McDowell, J.C., Green, R.F., Bechtold, J., Willner, S.P., Oey, M.S., Polonski, E., Cutri, R., 1994, *ApJS*, 95, 1
- Fabian, A.C., 1999, *MNRAS*, 308, L39
- Fanaroff, B.L., Riley, J.M., 1974, *MNRAS*, 167, 31
- Grandi, S.A., Osterbrock, D.E., 1978, *ApJ*, 220, 783
- Greene, J.E., Ho, L.C., 2005, *ApJ*, 630, 122
- Grupe, D., Mathur, S., 2004, *ApJ*, 606, L41
- Heckman T.M., Smith E.P., Baum S.A., van Breugel W.J.M., Miley G.K., Illingworth, G.D., Bothun, G.D., Balick, G.D., 1986, 311, 526
- Heiles, C., Kulkarni, S., 1988, In Kellerman, K., Verschuur, G., eds, *Galactic & Extragalactic Radio Astronomy*, Kluwer, Berlin, p95
- Holt, J., Tadhunter, C.N., Morganti, R., 2003, *MNRAS*, 342, 995
- Holt, J., 2005, PhD Thesis, University of Sheffield
- Hopkins, P.F., Hernquist, L., Cox, T.J., di Matteo, T., Martini, P., Robertson, B., Springel, V., 2005, *ApJ*, 630, 705
- Jauncey, D.L. & The SHEVE Team, 1994, in *Very high angular resolution imaging*, IAU158, eds J.G., Robertson & W.J. Tango (Dordrecht: Kluwer), p131
- Johnston, H.M., Hunstead, R.W., Cotter, G., Sadler, E.M., 2005, *MNRAS*, 356, 515
- Kauffmann, G., Haehnelt, M., 2000, *MNRAS*, 311, 576
- King, E., 1994, PhD thesis, University of Tasmania
- Madau, P., Ferguson, H.C., Dickinson, M.E., Giavalisco, M., Steidel, C.C., Fruchter, A., 1996, *MNRAS*, 283, 1388
- Maiolino, R., Marconi, A., Salvati, M., Risaliti, G., Severgnini, P., Oliva, E., La Franca, F., Vanzì, L., 2001, *A&A*, 365, 28
- Mathur, S., 2000, *MNRAS*, 314, L17
- Mathur, S., Grupe, D., 2005, *A&A*, 432, 436
- McLure, R.J., Dunlop, J.S., 2002, *MNRAS*, 331, 795
- Mihos, C.J., Hernquist, L., 1996, 464, 641
- Morganti, R., Robinson, A., Fosbury, R.A.E., di Serego Alighieri, S., Tadhunter, C.N., Malin, D.F., 1991, *MNRAS*, 249, 91
- Morganti, R., Oosterloo, T., Tadhunter, C.N., 2001, *MNRAS*, 323, 331
- Morganti, R., Oosterloo, T., Emonts, B.H.C., van der Hulst, J.M., Tadhunter, C.N., 2003, *ApJ*, 593, L69
- Morganti, R., Oosterloo, T.A., Tadhunter, C.N., van Moorsel, G., Emonts, B., 2005a, *A&A*, 439, 521
- Morganti, R., Tadhunter, C.N., Oosterloo, T.A., 2005b, *A&A*, 444, L9
- Morganti, R., Tadhunter, C., Oosterloo, T., 2005, *A&ALett*, in press
- Osterbrock, D.E., Pogge, R.W., 1985, *ApJ*, 297, 166
- Osterbrock, D.E., 1987, *Lecture Notes in Physics*, 307, 1
- Osterbrock, D.E., 1989, *The Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*, University Science Books
- Pounds, K., Done, C., Osborne, J., 1995, *MNRAS*, 277, L5
- Preston, R.A., The SHEVE Team, 1993, In: *Sub-arcsecond radio astronomy*, R.J. Davis & R.S. Booth (eds), CUP, p428
- Punsly, B., Tingay, S.J., 2005, *ApJ*, 633, L93
- Rieke, G.H., Lebofsky, M.J., 1985, *ApJ*, 288, 618
- Robinson T. G., Tadhunter C. N., Axon D. J., Robinson A., 2000, *MNRAS*, 317, 922
- Rupke, D.S., Veilleux, S., Sanders, D.B., 2005a, *ApJ*, 632, 751
- Rupke, D.S., Veilleux, S., Sanders, D.B., 2005b, *ApJS*, 160, 115
- Sanders, D.B., Mirabel, I.F., 1996, *ARA&A*, 34, 1996
- Sault, R.J., Teuben, P.J., Wright, M.C.H., 1995, in Shaw R., Payne, H.E., Hayes, J.J.E., eds, *ASP Conf. Ser. Vol. 11, Astronomical Data Analysis Software & Systems IV*, Astron. Soc. Pac. San Francisco, P433
- Schlegel, D.J., Finkbeiner, D.P., Davis, M., 1998, *ApJ*, 500, 525
- Seaton, M.J., 1979, *MNRAS*, 187, 73
- Shaw M., Tadhunter C., Dickson R., Morganti R., 1995, *MNRAS*, 275, 703
- Shepherd, M.C., Pearson, T.J., Taylor, G.B., 1994, *BAAS*, 26, 987
- Silk, J., Rees, M.J., 1998, *A&A*, 331, L1
- Simpson, C., Rawlings, S., 2000, *MNRAS*, 317, 1023
- Smith, E.P., Heckman, T.M., 1989, *ApJ*, 341, 658
- Solorzano-Innarea, C., 2001, PhD thesis, University of Sheffield

- Storchi-Bergmann, T., Wilson, A.S., Mulchaey, J.S., Binette, L., 1996, *A&A*, 312, 357
- Tadhunter, C.N., Fosbury, R.A.E., Quinn, P.J., 1989, *MNRAS*, 240, 225
- Tadhunter, C., Scarrott S., Draper P., Rolph C., 1992, *MNRAS*, 256, 53p
- Tadhunter, C.N., Metz, S., Robinson, A., 1994, *MNRAS*, 268, 989
- Tadhunter, C.N., Dickson, R.C., Shaw, M.A., 1996, *MNRAS*, 281, 591
- Tadhunter, C.N., Wills, K., Morganti, R., Oosterloo, T., Dickson, R., 2001, *MNRAS*, 327, 227
- Tadhunter, C., Dickson, R., Morganti, R., Robinson, T.G., Wills, K., Villar-Martin, M., Hughes, M., 2002, *MNRAS*, 300, 977
- Tadhunter, C., Marconi, A., Axon, D., Wills, K., Robinson, T.G., Jackson, N., 2003, *MNRAS*, 342, 861
- Tadhunter, C., Robinson, T.G., Gonzalez Delgado, R.M., Wills, K., Morganti, R., 2005, *MNRAS*, 356, 480
- Taylor, M.D., Tadhunter, C.N., Robinson, T.G., 2003, *MNRAS*, 342, 995
- Tremaine, S., et al., 2002, *ApJ*, 574, 740
- Tzioumis, A., et al., 2002, *A&A*, 392, 841
- Veilleux, S., Cecil, G., Bland-Hawthorn, J., 2005, *ARA&A*, 43, 769
- Veron-Cetty, M.-P., Veron, P., Goncalves, A.C., 2001, *A&A*, 372, 730
- Villar-Martin, M., Tadhunter, C.N., Morganti, R., Clark, N., Kilean, N., Axon, D., 1998, *A&A*, 332, 479
- Villar-Martin, M., Tadhunter, C.N., Morganti, R., Axon, D.J., Koekemoer, A., 1999, *MNRAS*, 307, 24
- Wills, K.A., Tadhunter, C.N., Robinson, T.G., Morganti, R., 2002, *MNRAS*, 333, 221
- Wills, K.A., Morganti, R., Tadhunter, C.N., Robinson, T.G., Villar-Martin, M., 2004, *MNRAS*, 347, 771
- Zamanov, R., Marziani, P., Sulentic, J.W., Calvani, M., Dultzin-Hacyan, D., Bachev, R., 2002, *ApJ*, 576, L9